BEREICE BEREIC



ORGANO-MINERAL FERTILIZER NPK PHYTOSTIMULANT OBTAINED BY REACTION



GRENA STARTER 12.5.6 is recommended for its starter effect in spring fertilisations



SOURCE

Organic: meatmeal Mineral: ammonium sulphate, diammonium (DAP), urea, potassium chloride

Leucine

Physical state: micro 2 mm - pellet 4 mm

Packaging available: 25 kg bags - 500 kg bags GRENA STARTER 12.5.6 is obtained by reaction between the GRENA organic matrix and high-quality mineral materials.

The high percentage of total nitrogen is harmonized by a gradual presence of ammonia, ureic and organic nitrogen, from which an ideal release curve for spring growth is obtained. **The simultaneous presence of phosphorus and potassium creates a synergic nutritive** action on the root system, making the plant treated with GRENA STRARTED 12.5.6 stronger and healthier.

In fact, nitrogen stimulates a more lush bloom and the formation of more tender vegetable tissue, while potassium stimulates the formation of thicker cell walls, which increase the intrinsic resistance of the tissue to parasitic infections.

GRENA STARTER in micro form **allows a significant reduction in doses** and allows a homogeneous and practically invisible distribution, an important feature when the product is used on ornamental lawns, or on golf courses and football fields.

AMINO ACIDS IN GREN	A MATRIX
Aspartic Acid	1.25 g/100 g
Glutamic Acid	1.62 g/100 g
Alanine	1.02 g/100 g
Arginine	0.83 g/100 g
Phenylalanine	0.56 g/100 g
Glycine	0.95 g/100 g
Hydroxyproline	0.22 g/100 g
Isoleucine	0.62 g/100 g
Histidine	0.31 g/100 g
Leucine	1.10 g/100 g
Lysine	0.56 g/100 g
Proline	0.85 g/100 g
Serine	0.87 g/100 g
Tyrosine	0.33 g/100 g
Threonine	0.59 g/100 g
Valine	0.80 g/100 g
Cysteine and Cystine	0.18 g/100 g
Methionine	0.19 g/100 g
Tryptophan	0.09 g/100 g
FREE AMINO ACIDS	
Glutamic Acid	0.06 g/100 g
Alanine	0.12 g/100 g

MICRO-ELEMENTS

В	2.30 mg/kg
Fe	330 mg/kg
Mn	18.6 mg/kg
Cu	2.87 mg/kg
Zn	33.6 mg/kg

COMPOSITION	
Organic matter	40%
Organic substance (Cx1.724)	26%
Amino acids and proteins (Nx6.25)	10%
Humic and fulvic acids	6.9%
Humidity	7%
Total nitrogen (N)	12%
Organic nitrogen (N)	1%
Ammoniacal nitrogen (N)	10%
Ureic nitrogen (N)	1%
Phosphoric anhydride (P ₂ O ₅)	5%
Potassium oxide (K ₂ O) soluble in water	6%
Organic carbon (C)	15%
Sulphuric anhydride (SO3)	11%
Calcium (CaO) natural origin	8%
C/N	1.25
Specific weight	0.85 kg/L

CROP	TIMING*	APPLICATION [*]	DOSAGE/HA*
Vineyards and orchards	autumn - winter	localized distribution per row	500-700 kg/ha
Olive groves	autumn - winter	localized distribution per row	1400-1600 kg/ha
Citrus fruits	autumn - spring	scatter the product in a dosage of 3/4 kg per plant	1400-1600 kg/ha
Open field crops	pre-sowing or pre-transplant	scatter the product in soil preparation	600-800 kg/ha
Ornamental lawns and gardens	winter - spring	scatter the product in soil	800-1000 kg/ha

0.05 g/100 g

*guidelines only, for the correct use of our products, please consult a specialist.